



What is Mytesi?

Mytesi is a prescription medicine that helps relieve symptoms of diarrhea not caused by an infection in adults living with HIV/AIDS on antiretroviral therapy (ART).

HOW TO TAKE MYTESI

1 Mytesi 125 mg
DELAYED-RELEASE
TABLET IS TAKEN **2 TIMES**
EACH DAY

YOU CAN TAKE IT
WITH OR WITHOUT
FOOD



Mytesi tablets should
NOT be crushed
or chewed

Tablets should be
swallowed whole



NapoCares™ Patient Support
Programs can help you start and
continue your Mytesi® treatment.

To enroll in NapoCares or to learn
more about its programs, please
call **(888) 527-NAPO (6276)**
Monday-Friday 8 am-5 pm CST.

*All NapoCares programs are free of
charge for Mytesi patients.*



Copay Savings Program to help you
save on your out-of-pocket costs†



Patient Assistance Program
for uninsured patients*



Support to help you **understand
your insurance coverage**



Support for **Prior Authorizations
and Appeals**



Mail order pharmacy support



Patient call center to answer any
questions



Linkage to specialty pharmacies
to fill your prescriptions

Important Safety Information

Mytesi (crofelemer) is not approved to treat infectious diarrhea (diarrhea caused by bacteria, a virus, or a parasite). Before starting you on Mytesi, your healthcare provider will first be sure that you do not have infectious diarrhea. Otherwise, there is a risk you would not receive the right medicine and your infection could get worse. In clinical studies, the most common side effects that occurred more often than with placebo were upper respiratory tract (sinus, nose, and throat) infection (6%), bronchitis (4%), cough (4%), flatulence (3%), and increased bilirubin (3%).

See package insert for full prescribing information.

*Subject to eligibility. Restrictions apply. Rx only. Subject to change or cancellation at any time and without notice.

†Patient must be commercially insured for eligibility. Subject to change or cancellation at any time and without notice.

TRACKING YOUR PROGRESS

When taking Mytesi daily, most patients continue to see improvement in their symptoms even up to 20 weeks after starting treatment. It's important to continue taking Mytesi daily to see these improvements, even if the symptom relief seems gradual.

WHAT'S YOUR NUMBER?

Adapted from the
Bristol Stool Form Scale

CONSTIPATION

1



Separate hard lumps, like nuts

2



Lumpy and sausage-like

NORMAL

3



A sausage shape with cracks in the surface

4



Like a smooth, soft sausage or snake

DIARRHEA (LOOSE/WATERY STOOLS)

5



Soft blobs with clear-cut edges

6



Mushy consistency with ragged edges

7



Liquid consistency with no solid pieces

Date	Number of Loose/Watery Stools (5, 6, or 7 from above)	Notes
BASELINE before starting Mytesi Date: ____ / ____ / ____		
1 WEEK after starting Mytesi Date: ____ / ____ / ____		
4 WEEKS after starting Mytesi Date: ____ / ____ / ____		
8 WEEKS after starting Mytesi Date: ____ / ____ / ____		
20 WEEKS after starting Mytesi Date: ____ / ____ / ____		

NapoCares has support for Mytesi patients at no cost to them, whether they are uninsured, underinsured, or insured. To enroll in NapoCares or to learn more about its programs, please call 1-888-527-NAPO (6276) Monday-Friday 8 am - 5 pm CST.

Important Safety Information on the other side. See package insert for full prescribing information.